



Hugo B. Fischer Award Presentation

Michael L. MacWilliams, PhD, PE
April 17, 2023




Thank You!

- J. Keith Rigby
- Robert Street
- Peter Kitanidis
- Vincenzo Casulli
- Ed Gross
- Ralph Cheng
- Wim Kimmerer
- John DeGeorge
- Richard Rachiele
- Steve Monismith
- Frank Wu
- Bruce Herbold
- Aaron Bever
- Ted Sommer
- UnTRIM User Group
- Data collectors and data repositories
- Many other clients and collaborators
- My family



Twenty Years of Delta Model Advancements 2003–2005 TRIM3D



Simulating Periodic Stratification in the San Francisco Estuary

Edward S. Gross,¹ Michael L. MacWilliams² and Wim Kimmerer³

Abstract

Three-dimensional simulations of circulation in the San Francisco Estuary were performed with the three-dimensional hydrodynamic model, TRIM3D, using a generic length scale turbulence closure model. The model was calibrated to reproduce observed tidal elevations and tidal currents in the San Francisco Estuary and then was applied during a period of intensive data collection. The model predicts tidal currents accurately and realistically simulates variability in salinity at both the seasonal and tidal time scale. The model results are consistent with the current conceptual understanding of stratification in the San Francisco Estuary and approximately predict the magnitude and tidal phasing of observed stratification. The simulation results were analyzed to improve understanding of periodic stratification. During spring tides, compression of salinity gradients near high water and weak tidal currents at and following high water allow the formation of weak and transient stratification. During neap tides stronger stratification is observed and predicted. This stratification is strongest during ebb when tidal straining is effective in creating stratification. Vertical turbulent mixing may be adequate to diminish vertical stratification during strong ebb tides while stratification can persist or increase through weak ebb tides.

Introduction

The hydrodynamic modeling effort presented is part of a larger effort to better understand the effects of freshwater inflow on the abundance of estuarine biota in the San Francisco Estuary (defined as the body of water from the Delta to the Golden Gate, including South San Francisco Bay, Central San Francisco Bay, San Pablo Bay, Carquinez Strait, Suisun Bay and the Sacramento-San Joaquin Delta). Freshwater flow into the San Francisco Estuary is regulated in part using a salinity standard defined as the position of 2 psu (practical salinity units) below bottom salinity. The location of 2 psu below salinity, known as X_2 , is reported as distance along the channel in kilometers from the Golden Gate (Jassby et al. 1995). The X_2 standard was developed based on observations that the survival and abundance of several estuarine fish species correlate with X_2 (Jassby et al. 1995). Mechanisms behind these correlations are unknown, but for some species they may be linked to the timing, location, and frequency of stratification in the estuary (Kimmerer 2004). We are using the TRIM3D model to investigate plausible mechanisms involving stratification, and must therefore ensure that the prediction of stratification by the model is realistic.

¹ Environmental Consultant, 1777 Spence Street, Berkeley, CA, 94709, ed.gross@baymodeling.com
² Environmental Consultant, P.O. Box 223174, San Francisco, CA 94112, michael@rivermodeling.com
³ Romberg Tiburon Center, 3152 Paradise Drive, Tiburon, CA 94920, kimmerer@tbc.edu

155

Copyright ASCE 2006

Estuarine and Coastal Modeling 2005

Source: Gross et al. 2005

Estuaries and Coasts
 DOI 10.1007/s12237-008-9124-x

Is the Response of Estuarine Nekton to Freshwater Flow in the San Francisco Estuary Explained by Variation in Habitat Volume?

Wim J. Kimmerer · Edward S. Gross · Michael L. MacWilliams

Received: 27 June 2008 / Revised: 13 November 2008 / Accepted: 23 November 2008
 © Coastal and Estuarine Research Federation 2009

Abstract Abundance of estuarine biota can vary with freshwater inflow through several mechanisms. One proposed mechanism is that the extent of physical habitat for an estuarine species increases with flow. We estimated the contribution of variation in habitat volume to the response of eight species of estuarine nekton to changes in freshwater flow in the San Francisco Estuary. Species selection functions for salinity and depth were developed for each species and for five additional species using frequency data sets. The TRIM3D hydrodynamic model was run for five study flow scenarios to determine volume by salinity and depth, and retention selection functions were used as a weighting factor to calculate an index of total habitat index. We then correlated total depth of abundance vs. flow for only two of the species examined. Therefore, other mechanisms must underlie responses of abundance to flow for most species.

Keywords Fish · Habitat · Freshwater flow · Retention selection function · San Francisco Estuary

W. J. Kimmerer (✉)
 Romberg Tiburon Center, San Francisco State University,
 1635 Paradise Drive,
 Tiburon, CA 94920, USA
 e-mail: wkimmerer@sfsu.edu

E. S. Gross
 4012 Tilden Road,
 Oakland, CA 94612, USA

M. L. MacWilliams
 P.O. Box 223174, San Francisco, CA 94112-3174, USA

Published online: 18 January 2009




Introduction
 Variability in freshwater flow is the principal mode of interannual and seasonal variation of physical conditions in many estuaries (Chandler 1964). River discharge mechanisms may be sensitive to climate change and increasing human control (Vitousek et al. 2000; Savari et al. 2003). Thus, understanding mechanisms by which estuarine ecosystems respond to freshwater flow should yield important insights into the dynamics of these ecosystems and their sensitivity to perturbation.

Biological populations in estuaries often vary with freshwater flow. Greater flow effects have been reported for phytoplankton production (Hagy 1993; Mallin et al. 1995; Shi et al. 1996) and for abundance or biomass of benthic invertebrates (Abram 1977; Kimmerer 1992; Mangun and Kelle 1992; Wilce 1992, 1994; Frazee et al. 2007) and fish densities (Fry 1977; Hsieh and Rutherford 1993; Jassby et al. 1995). Negative effects on biological populations on the river flow and biomass 1995, i.e., through effects of wastewater or pesticide stress (Deegan 1990; Kimmel and Adams 1992).

Various potential mechanisms have been proposed for positive effects of freshwater flow on biological populations (e.g., Niemi et al. 1980; Chen 1991; DeBusk and Fausch 1994; Kimmerer 2004, 05). One proposed mechanism is the increase in area or volume of physical habitat for biota that accompanies increases in freshwater flow (Kimmerer et al. 1995; Kimmerer 2004). This mechanism may explain increases in the abundance of Sacramento splittail, *Pogonias cromis* macrolepax, with freshwater flow in the upper San Francisco estuary (Chen et al. 1997). When high flow inundates floodplains adjacent to the estuary, splittail gain access to large areas of habitat, particularly for foraging and spawning (Fryer et al. 2004).

Source: Kimmerer et al. 2009

Twenty Years of Delta Model Advancements 2004–2007 UnTRIM



THREE-DIMENSIONAL HYDRODYNAMIC MODELING OF SAN PABLO BAY ON AN UNSTRUCTURED GRID

Michael L. MacWilliams¹ and Ralph T. Cheng²

ABSTRACT

A three-dimensional hydrodynamic model of San Francisco Bay was developed using the three-dimensional hydrodynamic model UnTRIM. The model was calibrated using continuous water level measurements and ADCIRC data in San Francisco Bay, and validated during an additional simulation period using current velocity measurements. The model was developed to support the Hamilton Wetlands Restoration Project, a joint undertaking by the U.S. Army Corps of Engineers and the California Coastal Conservancy to restore 2.6 km² of tidal marsh bordering San Pablo Bay. The restoration effort is expected to make use of more than 8.1 million m³ of dredged materials to raise the elevation of isolated wetlands. The placement of an Aquatic Transfer Facility (ATF) in San Pablo Bay is being considered by the U.S. Army Corps of Engineers to serve as a temporary holding site for dredge sediments before they are transferred to the Hamilton Wetlands restoration site. The San Francisco Bay model presented in this paper was developed as part of a larger study to evaluate potential impacts of the proposed ATF on circulation and sediment transport dynamics in San Pablo Bay. This paper presents the model calibration and validation, while the full analysis of proposed ATF conditions is presented in a separate technical report.

1. INTRODUCTION

A three-dimensional hydrodynamic model of San Francisco Bay was developed as part of the Hamilton Wetlands Restoration Project, a joint undertaking by the U.S. Army Corps of Engineers and the California Coastal Conservancy. It is estimated that 85 to 90 percent of the historic tidal marshes bordering San Francisco Bay have been filled or significantly altered over the past two centuries (SBSF, 2006). The Hamilton Wetlands Restoration Project is part of a growing effort to restore a portion of these former marshes to tidal action. The Hamilton Wetlands Restoration site, historically dominated by tidal salt marsh habitat, was converted first to agricultural and then for use as the Hamilton Army Airfield (USACE, 1985). Since the site was originally established, it has subsided six to nine feet below mean sea level, and more than 8.1 million m³ of dredged materials are expected to be used to raise the elevations of these isolated wetlands (USACE, 1985). In order to facilitate the transfer of dredged material to the project site, and Aquatic Transfer Facility (ATF) is being considered by the U.S. Army Corps of Engineers to serve as a temporary holding site for dredge sediments before they are transferred to the Hamilton Wetlands Restoration site. The basic concept of the ATF consists of an excavated basin located in relatively deep water which can be

¹ Environment Consultant, P.O. Box 221174, San Francisco, CA 94122-1174, USA (mwilliams@environmental.com)
² Project Chief, Estuarine Hydrodynamics, U.S. Geological Survey, Menlo Park, CA 94025 (rcheng@usgs.gov)

Source: MacWilliams and Cheng 2006

THREE-DIMENSIONAL HYDRODYNAMIC MODELING OF THE SAN FRANCISCO ESTUARY ON AN UNSTRUCTURED GRID

Michael L. MacWilliams¹, Edward S. Gray²,
John F. DeGeorge³, Richard R. Rachate⁴

¹ Environment Consultant, P.O. Box 221174, San Francisco, CA 94122, mwilliams@environmental.com
² Environment Consultant, 4617 Paper Street, Oakland, CA 94612, edgray@environmental.com
³ Resource Management Associates, 4171 Sausal Valley Road, Suite 1, Foothill, CA 94134, jdegeorge@rma.com
⁴ Resource Management Associates, 4171 Sausal Valley Road, Suite 1, Foothill, CA 94134, rrachate@rma.com

ABSTRACT

Three-dimensional simulations of circulation in the San Francisco Estuary were performed with the three-dimensional unstructured grid hydrodynamic model UnTRIM. The model was developed to support the Delta Risk Management Strategy (DRMS) funded by the California Department of Water Resources. The model applications build on previous TRIM and UnTRIM applications. A model grid consisting of quadrilaterals and triangles was developed that extends from the Pacific Ocean through San Francisco Bay and farther into the Sacramento-San Joaquin Delta than previous TRIM and UnTRIM applications. In addition, a state-of-the-art turbulence closure was incorporated into the UnTRIM model from the TRIM model. This paper describes a portion of the hydrodynamic and salinity calibration of the resulting San Francisco Estuary UnTRIM model.

Keywords: three-dimensional, unstructured grid, primary, salinity, UnTRIM, San Francisco Bay, Sacramento-San Joaquin Delta, turbulence closure, generic length scale, calibration



1 INTRODUCTION

The Sacramento-San Joaquin Delta is a critical resource to the state of California because the Delta is a source of drinking water for roughly 2 out of 3 Californians. However, the 2,800 km² of islands in the Delta region are at risk of inundation from levee failures. These deeply subsided islands are protected by levees typically 4 to 5 meters high which, in most cases, are engineered levees and are constructed partially with peat and other weak and compressible soils. The Delta Risk Management Study has been funded by the California Department of Water Resources (CDWR) to look at vulnerability of the Delta and assess major risks to the Delta resources from floods, seepage, subsidence, and earthquakes. Part of this effort involves the application of hydrodynamic models to estimate the effects of levee failures on salinity in the Delta. Levee failures in the Delta generally result in increased salinity as islands flood and brackish water from Suisun Bay is entrained into the Delta. Increased salinity can result in exceedance of water quality objectives for drinking water causing interruption of water exports, resulting in a large economic impact.


Several hydrodynamic and water quality simulation tools are applied in the DRMS project, ranging from a tidally-averaged advection-dispersion model, which can perform a year of salinity projections in 1 minute of computer time, to the sophisticated and computationally intensive three-dimensional model described here. The first phase of the DRMS work involves quantification of risk and consequences of Delta levee failures while the second phase will evaluate risk reduction actions that can be taken to reduce risks of levee failures and the impacts of levee failures.

Source: MacWilliams et al. 2007

Twenty Years of Delta Model Advancements 2008–2009 UnTRIM




Source: MacWilliams et al. 2008



Source: MacWilliams and Gross 2010

Twenty Years of Delta Model Advancements 2010–2023 UnTRIM



SAN FRANCISCO ESTUARY & WATERSHED SCIENCE

Three-Dimensional Modeling of Hydrodynamics and Salinity in the San Francisco Estuary: An Evaluation of Model Accuracy, X2, and the Low-Salinity Zone

Michael L. MacWilliams^{1*}, Aaron J. Bever¹, Edward S. Gross², Gerard S. Ketefian², and Wim J. Kimmerer³

Volume 13, Issue 1 | April 2015
doi: <http://dx.doi.org/10.1544/estw.2015.13.iss1.w12>
* Corresponding author: micwill@deltaonline.org

¹ Delta Modeling Associates, Inc., 870 Market Street, Suite 1074, San Francisco, CA 94102 USA
² Resource Management Associates, Inc., Berkeley, CA 94704 USA
³ Romberg Tiburon Center, San Francisco State University, Tiburon CA 94920 USA

ABSTRACT


The three-dimensional UnTRIM San Francisco Bay-Delta model was applied to simulate tidal hydrodynamics and salinity in the San Francisco Estuary (estuary) using an unstructured grid. We compared model predictions to observations of water level, tidal flow, current speed, and salinity collected at 137 locations throughout the estuary. A quantitative approach based on multiple model assessment metrics was used to evaluate the model's accuracy for each comparison. These comparisons demonstrate that the model accurately predicted water level, tidal flow, and salinity during a 3-year simulation period that spanned a large range of flow and salinity conditions. The model is therefore suitable for detailed investigation of circulation patterns and salinity distributions in the estuary.

The model was used to investigate the location, and spatial and temporal extent of the low-salinity zone (LSZ), defined by salinity between 0.5 and 6 psu. We calculated X2, the distance up the axis of the estuary to the daily-averaged 2-psu near-bed salinity, and the spatial extent of the LSZ for each day during the 3-year simulation. The location, area, volume, and average depth of the low-salinity zone varied with X2; however this variation was not monotonic and was largely controlled by the geometry of the estuary.

We used predicted daily X2 values and the corresponding daily Delta outflow for each day during the 3-year simulation to develop a new equation to relate X2 to Delta outflow. This equation provides a conceptual improvement over previous equations by allowing the time constant for daily changes in X2 to vary with flow conditions. This improvement resulted in a smaller average error in X2 prediction than previous equations. These analyses demonstrate that a well-calibrated three-dimensional (3-D) hydrodynamic model is a valuable tool for investigating the salinity distributions in the estuary, and their influence on the distribution and abundance of physical habitat.

Source: MacWilliams et al. 2015

Sediment Transport and Morphology



Water Geology 44(3):205-231
Contents lists available at ScienceDirect

Marine Geology

Journal of ScienceDirect www.elsevier.com/locate/margeo

Simulating sediment transport processes in San Pablo Bay using coupled hydrodynamic, wave, and sediment transport models

Aaron J. Bever^a, Michael L. MacWilliams^a

^a Delta Modeling Systems Inc., 1000 Redwood, Suite 300, San Francisco, CA 94132, United States

ARTICLE INFO
 Article history:
 Received 10 March 2013
 Received in revised form 27 June 2013
 Accepted 12 July 2013
 Available online 23 August 2013

Keywords:
 San Pablo Bay
 Sediment transport
 Hydrodynamics
 SWAN
 SediMorph

ABSTRACT
 San Pablo Bay is the only sub-estuary of the San Francisco Estuary which is dominated by the tidal delta. It has been identified as a high priority area for management of the estuary. To improve our understanding of the bay, we developed a hydrodynamic, wave, and sediment transport model using the coupled SWAN, Hydrodynamics, and SediMorph models. The model was used to simulate the bay's hydrodynamics, sediment transport, and morphology over a 10-year period. The model results show that the bay is highly dynamic and that sediment transport is dominated by the tidal delta. The model results also show that the bay is a highly dynamic system and that sediment transport is dominated by the tidal delta. The model results show that the bay is highly dynamic and that sediment transport is dominated by the tidal delta. The model results also show that the bay is a highly dynamic system and that sediment transport is dominated by the tidal delta.

© 2013 Elsevier B.V. All rights reserved.

Source: Bever and MacWilliams 2013

Estuaries and Coasts 2010, 41: 1942–1957
<https://doi.org/10.1007/s12237-010-9184-5>

Influence of an Observed Decadal Decline in Wind Speed on Turbidity in the San Francisco Estuary

Aaron J. Bever^a, Michael L. MacWilliams^a, David K. Fulkerson^a

^a Delta Modeling Systems Inc., 1000 Redwood, Suite 300, San Francisco, CA 94132, United States

Received 20 June 2010 / Revised 22 March 2011 / Accepted 1 April 2011 / Published online 15 April 2011
 © The Author(s) 2011

ABSTRACT
 Turbidity is an important habitat component in estuaries for many fishes and affects a range of other ecological functions. Decadal-scale declines in turbidity have been observed in the San Francisco Estuary (SFE) with Redfish generally unaffected by reductions in turbidity, but a variety of other species, including juvenile salmonids, have experienced significant declines. We analyzed turbidity data from 1995 through 2007 and found statistically significant declines of 17% over wind speed around the Estuary. This wind speed reduction was associated with a decline in wind speed, which was in turn associated with a decline in wind speed around the Estuary. The reduction in wind speed over the past 20 years was projected to result in a decrease in turbidity of 14% to 25% in SFEs. These findings highlight the need to consider the effects of observed declines in wind speed on sediment supply over the past 20 years, have resulted in reduced turbidity in the San Francisco Estuary from October through January. This decline in turbidity in SFEs may potentially have negative effects on habitat for fish. In the development of fish habitat, which are more consistently caught in relatively turbid waters.





KEYWORDS
 Turbidity · Delta · Sediment · Hydrodynamic modeling · Sediment transport · San Francisco Bay · Climate change

Introduction
 Motivation and Background
 Water column turbidity is an important habitat component in many estuarine systems, and changes in turbidity can have significant management implications. In the San Francisco Estuary (SFE), reduced turbidity has been observed and has been linked to a decline in fish populations (Fulton et al. 2007; Sarmiento and Hagan 2007), and, in the past, phytoplankton growth was light-limited (Cham 1971; Aguirre and Cham 1981). Deltaic turbidity distribution and turbidity variability are well documented for areas ranging from California water supply for drinking, fisheries and channels. Biochemists in

Source: Bever et al. 2018

Standards for Assessment of Model Accuracy

Model accuracy		Water level	Flow	Salinity	Current speed
Skill accuracy	Accurate	>0.975	>0.975	>0.85	>0.9
	Acceptable	0.95 – 0.975	0.95 – 0.975	0.7 – 0.85	0.8 – 0.9
	Poor agreement	<0.95	<0.95	<0.7	<0.8
Target accuracy	Very accurate	0.0 - 0.25			
	Accurate	0.25 - 0.5			
	Acceptable	0.5 - 1.0			
	Poor agreement	> 1.0			



APRIL 2015

SAN FRANCISCO ESTUARY & WATERSHED SCIENCE

Three-Dimensional Modeling of Hydrodynamics and Salinity in the San Francisco Estuary: An Evaluation of Model Accuracy, X2, and the Low-Salinity Zone

Michael L. MacWilliams¹, Aaron J. Bever¹, Edward S. Gross², Gerard S. Ketefian², and Win J. Kimmerer³

Volume 13, Issue 1 | April 2015
doi: <http://dx.doi.org/10.154479/estw.2015v13i01a012>
Corresponding author: mike@slab.wisc.edu

¹ Delta Modeling Associates, Inc., 670 Market Street, Suite 1074, San Francisco, CA 94102 USA
² Resource Management Associates, Inc., Berkeley, CA 94704 USA
³ Reenberg Tiburon Center, San Francisco State University, Tiburon CA 94920 USA

ABSTRACT



The three-dimensional UnTRIM San Francisco Bay-Delta model was applied to simulate tidal hydrodynamics and salinity in the San Francisco Estuary (estuary) using an unstructured grid. We compared model predictions to observations of water level, tidal flow, current speed, and salinity collected at 137 locations throughout the estuary. A quantitative approach based on multiple model assessment metrics was used to evaluate the model's accuracy for each comparison. These comparisons demonstrate that the model accurately predicted water level, tidal flow, and salinity during a 3-year simulation period that spanned a large range of flow and salinity conditions. The model is therefore suitable for detailed investigation of circulation patterns and salinity distributions in the estuary.

The model was used to investigate the location, and spatial and temporal extent of the low-salinity zone (LSZ), defined by salinity between 0.5 and 6 psu. We calculated X2, the distance up the axis of the estuary to the daily-averaged 2-psu near-bed salinity, and the spatial extent of the LSZ for each day during the 3-year simulation. The location, area, volume, and average depth of the low-salinity zone varied with X2; however this variation was not monotonic and was largely controlled by the geometry of the estuary.


We used predicted daily X2 values and the corresponding daily Delta outflow for each day during the 3-year simulation to develop a new equation to relate X2 to Delta outflow. This equation provides a conceptual improvement over previous equations by allowing the time constant for daily changes in X2 to vary with flow conditions. This improvement resulted in a smaller average error in X2 prediction than previous equations. These analyses demonstrate that a well-calibrated three-dimensional (3-D) hydrodynamic model is a valuable tool for investigating the salinity distributions in the estuary, and their influence on the distribution and abundance of physical habitat.

Source: MacWilliams et al. 2015

X2 and the Low Salinity Zone




Source: Bever et al. 2016





Source: Delta Modeling Associates 2014

Subgrid Bathymetry



Seasonally Averaged Salinity Gradient -- January to June



RESEARCH

3-D Simulations of the San Francisco Estuary with Subgrid Bathymetry to Explore Long-Term Trends in Salinity Distribution and Fish Abundance

Michael L. MacWilliams¹, Aaron J. Bever¹, and Erin Foresman²

Volume 14, Issue 2 | Article 3
doi: <http://dx.doi.org/10.15447/sfwes.2016v14a2a3>

* Corresponding author: mwilliams@ancher.org
¹ Ancher OEA, LLC
 San Francisco, CA 94111 USA
² U.S. Environmental Protection Agency
 San Francisco, CA 94105 USA

ABSTRACT

The UnTRIM hydrodynamic model was applied to San Francisco Bay and the Sacramento-San Joaquin Delta (Delta) using a coarse-resolution model grid with bathymetry represented at a finer subgrid scale. We simulated a 35-year period, spanning from January 1, 1980 through December 31, 2014. This simulation was used to develop salinity distribution maps to facilitate visualization of fish distribution and abundance data. We compared predicted salinity from the coarse-grid UnTRIM Bay-Delta model to continuous salinity monitoring observations as well to the measured surface salinity from San Pablo Bay through the Delta at a total of 5,542 times and locations where surface salinity was observed as part of several long-term fish monitoring programs: the Fall Midwater Trawl, Summer Trawl Survey, and San Francisco Bay Study. The coarse-grid UnTRIM Bay-Delta model was shown to accurately predict hydrodynamics and the spatial distribution of salinity over both a 3-year detailed validation period and over the full 35-year analysis period. The predicted

salinity was used to calculate the daily position of X2 and the daily-averaged area of the Low Salinity Zone (LSZ) for each day during the 35-year simulation. Our analysis highlights the influence of multi-year climate patterns, shorter-duration weather patterns, and Delta outflow on salinity distribution. We used the predicted salinity to develop maps of salinity distribution over seven periods for six fish species, and combined the salinity maps with historic fish sampling data to allow for visualization of fish abundance and distribution for 33 years between 1980 and 2012. These maps can be used to explore how different species respond to annual differences in salinity distributions in the San Francisco Estuary, and to expand the understanding of the relationships among salinity and fish abundance, distribution, and population resiliency.

KEY WORDS



San Francisco Bay, Hydrodynamic Modeling, UnTRIM, Low Salinity Zone, Fall Midwater Trawl, Bay Study, Fish Abundance, X2

INTRODUCTION

Long-term fisheries monitoring programs provide a valuable resource for understanding trends in fish abundance and distribution. These long-term monitoring programs in locations such as San

Source: MacWilliams et al. 2016a

Planning Large-Scale Flow Operations for Management of Estuarine Habitat



Percent time salinity is less than 6 PSU

Source: Sommer et al. 2020

PLOS ONE

RESEARCH ARTICLE

Evaluation of a large-scale flow manipulation to the upper San Francisco Estuary: Response of habitat conditions for an endangered native fish

Tad Sommer^{1,2}, Rosemary Hartman¹, Mistal Kozlov¹, Michael Kozlovich^{1,3}, J. Louise Conrad¹, Michael Hill-Forsell¹, Adam Brown¹, Christine Baruffi¹, April Hennessy¹, Michael Decker¹

¹ California Department of Water Resources, Sacramento, California, United States of America, ² Delta Science Program, Sacramento, California, United States of America, ³ Arroyo-CO2, LLC, San Francisco, California, United States of America, ⁴ California Department of Fish and Wildlife, Sacramento, California, United States of America, ⁵ U.S. Bureau of Reclamation, Sacramento, California, United States of America

tsommer@water.ca.gov

OPEN ACCESS

Abstract

While flow is known to be a major driver of estuarine acceptances, targeted flow manipulations are not because fish systems are extremely variable in space and time, and because the necessary infrastructure is rarely available. In summer 2018 we used a unique water control structure in the San Francisco Estuary (DTE) to direct a managed flow pulse into Suisun Marsh, one of the largest contiguous tidal wetlands on the west coast of the United States. The action was designed to increase habitat suitability for the endangered Delta Smelt (*Hypomesus transpacificus*) in a small estuary fish watershed in the upper DTE. The approach was to increase Sacramento River inflow (rivers) to about an additional 100 m³/s of pulse of the salinity water into Suisun Marsh during August, a critical time period for juvenile Delta Smelt rearing. Three-dimensional modeling showed that directing additional low salinity water into Suisun Marsh (Flow Action) substantially expanded the area of low salinity habitat by Delta Smelt that persisted beyond the period of DTE/COS operations. Field monitoring showed that both the ecological and water quality benefits in Suisun Marsh, representing better habitat conditions, than the upstream Sacramento River region throughout the study period. The Flow Action had no detectable effects on phytoplankton abundance, and did Suisun Marsh above average levels of these prey species in comparison to the Sacramento River. Field monitoring data suggested that steel head of Delta Smelt entered Suisun Marsh from the Sacramento River during the period of Flow Action. Comparison of the salinity effects of the Flow Action to historical catch data for Suisun Marsh further supported our hypothesis that the Flow Action would have been beneficial to the fish species. Our study provides insight into both the potential use of targeted flow manipulations to support endangered fishes such as Delta Smelt, and into the general response of estuarine habitat to flow management.

Introduction

While freshwater inflow has been a major focus of resource management in estuaries, including the upper San Francisco Estuary, there is a growing interest in using forward flow actions to maximize benefits for specific regions, habitats, and species. As a test of this concept, in summer 2018, we used a managed flow pulse to target an ecologically important region, a freshwater tidal slough complex (Cacho Slough Complex-CSC). Our goal was to improve estuarine habitat by increasing net flows through CSC to enhance downstream transport of lower trophic level resources, an important driver for fishes such as the endangered Delta Smelt (*Hypomesus transpacificus*). We used regional water infrastructure to direct 14.5 million m³ of Sacramento River flow into its adjacent Suisun Bay (Suisun Bay), where the pulse continued through CSC. Simulations using a 3-D hydrodynamic model (Delft3D) indicated that the managed flow pulse had a large effect on the flow of water through Suisun Marsh, and between CSC and further downstream. Multiple water quality concentrations (specific conductivity, dissolved oxygen, nutrients (NO₃⁻, NO₂⁻, NH₄⁺, PO₄³⁻) varied across the study region, and showed a strong response to the flow pulse. In addition, the lower Sacramento River had increased phytoplankton biomass and improved food quality indices (estimated from long-chain essential fatty acids) after the flow pulse. The managed flow pulse resulted in increases in quantities of zooplankton (zooplankton) demonstrating potential advection from upper floodplain channels into the target CSC and Sacramento River region. This study was conducted during a single year, which may have had unique characteristics; however, we believe that our study is an instructive example of how a relatively modest change in net flows can generate measurable changes in ecologically relevant metrics, and how an adaptive management action can help inform resource management.

KEY WORDS

Yolo Bypass, flow pulse, food quality, plankton, water quality, management action

Introduction

While freshwater inflow has been a major focus of resource management in estuaries, including the upper San Francisco Estuary, there is a growing interest in using forward flow actions to maximize benefits for specific regions, habitats, and species. As a test of this concept, in summer 2018, we used a managed flow pulse to target an ecologically important region, a freshwater tidal slough complex (Cacho Slough Complex-CSC). Our goal was to improve estuarine habitat by increasing net flows through CSC to enhance downstream transport of lower trophic level resources, an important driver for fishes such as the endangered Delta Smelt (*Hypomesus transpacificus*). We used regional water infrastructure to direct 14.5 million m³ of Sacramento River flow into its adjacent Suisun Bay (Suisun Bay), where the pulse continued through CSC. Simulations using a 3-D hydrodynamic model (Delft3D) indicated that the managed flow pulse had a large effect on the flow of water through Suisun Marsh, and between CSC and further downstream. Multiple water quality concentrations (specific conductivity, dissolved oxygen, nutrients (NO₃⁻, NO₂⁻, NH₄⁺, PO₄³⁻) varied across the study region, and showed a strong response to the flow pulse. In addition, the lower Sacramento River had increased phytoplankton biomass and improved food quality indices (estimated from long-chain essential fatty acids) after the flow pulse. The managed flow pulse resulted in increases in quantities of zooplankton (zooplankton) demonstrating potential advection from upper floodplain channels into the target CSC and Sacramento River region. This study was conducted during a single year, which may have had unique characteristics; however, we believe that our study is an instructive example of how a relatively modest change in net flows can generate measurable changes in ecologically relevant metrics, and how an adaptive management action can help inform resource management.

KEY WORDS

Yolo Bypass, flow pulse, food quality, plankton, water quality, management action

Introduction

While freshwater inflow has been a major focus of resource management in estuaries, including the upper San Francisco Estuary, there is a growing interest in using forward flow actions to maximize benefits for specific regions, habitats, and species. As a test of this concept, in summer 2018, we used a managed flow pulse to target an ecologically important region, a freshwater tidal slough complex (Cacho Slough Complex-CSC). Our goal was to improve estuarine habitat by increasing net flows through CSC to enhance downstream transport of lower trophic level resources, an important driver for fishes such as the endangered Delta Smelt (*Hypomesus transpacificus*). We used regional water infrastructure to direct 14.5 million m³ of Sacramento River flow into its adjacent Suisun Bay (Suisun Bay), where the pulse continued through CSC. Simulations using a 3-D hydrodynamic model (Delft3D) indicated that the managed flow pulse had a large effect on the flow of water through Suisun Marsh, and between CSC and further downstream. Multiple water quality concentrations (specific conductivity, dissolved oxygen, nutrients (NO₃⁻, NO₂⁻, NH₄⁺, PO₄³⁻) varied across the study region, and showed a strong response to the flow pulse. In addition, the lower Sacramento River had increased phytoplankton biomass and improved food quality indices (estimated from long-chain essential fatty acids) after the flow pulse. The managed flow pulse resulted in increases in quantities of zooplankton (zooplankton) demonstrating potential advection from upper floodplain channels into the target CSC and Sacramento River region. This study was conducted during a single year, which may have had unique characteristics; however, we believe that our study is an instructive example of how a relatively modest change in net flows can generate measurable changes in ecologically relevant metrics, and how an adaptive management action can help inform resource management.

KEY WORDS

Yolo Bypass, flow pulse, food quality, plankton, water quality, management action


Source: Sommer et al. 2020

Source: Frantzych et al. 2021


Closing Thoughts

“The challenge for the multi-dimensional modeler then becomes to take the enormous amount of information generated by the model and present it in a way that can be used to increase understanding of the system, without averaging out all of the important details.”

MacWilliams et al. 2016b (*State of Bay-Delta Science*)



Source: MacWilliams et al. 2016b



Source: MacWilliams et al. 2022

Thank you!



Source: MacWilliams et al. 2022

References

- Bever, A.J., and M.L. MacWilliams, 2013. "Simulating Sediment Transport Processes in San Pablo Bay Using Coupled Hydrodynamic, Wave, and Sediment Transport Models." *Marine Geology* 345:235–253. November 1, 2013.
- Bever, A.J., M.L. MacWilliams, B. Herbold, L.R. Brown, and F.V. Feyrer, 2016. "Linking Hydrodynamic Complexity to Delta Smelt (*Hypomesus transpacificus*) Distribution in the San Francisco Estuary, USA." *San Francisco Estuary and Watershed Science* 14(1):27. Available at: <https://doi.org/10.15447/sfews.2016v14iss1art3>.
- Bever, A.J., M.L. MacWilliams, and D.K. Fullerton, 2018. "Influence of an Observed Decadal Decline in Wind Speed on Turbidity in the San Francisco Estuary." *Estuaries and Coasts* 41:1943–1967. Available at: <https://doi.org/10.1007/s12237-018-0403-x>.
- Delta Modeling Associates (Delta Modeling Associates, Inc.), 2014. Low Salinity Flip Book. Version 2.0. December 31, 2014.
- Frantzich, J., B.E. Davis, M.L. MacWilliams, A.J. Bever, and T. Sommer, 2021. "Use of a Managed Flow Pulse as Food Web Support for Estuarine Habitat." *San Francisco Estuary and Watershed Science* 19(3). Available at: <https://doi.org/10.15447/sfews.2021v19iss3art3>.
- Gross, E.S., M.L. MacWilliams, and W. Kimmerer, 2005. *Simulating Periodic Stratification in San Francisco Bay*. Paper presented at the 9th International Conference on Estuarine and Coastal Modeling (Charleston, South Carolina); October–November 2005.

References (Continued)

Kimmerer, W.J., E.S. Gross, and M.L. MacWilliams, 2009. "Is the response of the Estuarine Nekton to Freshwater Flow in the San Francisco Estuary Explained by Variation in Habitat Volume?" *Estuaries and Coasts* 32:375–389. DOI: 10.1007/s12237-008-9124-x.

MacWilliams, M.L. and R.T. Cheng, 2006. *Three-Dimensional Hydrodynamic Modeling of San Pablo Bay on an Unstructured Grid*. 7th International Conference on Hydroscience and Engineering (Philadelphia, Pennsylvania); September 2006.

MacWilliams, M.L. and E.S. Gross, 2010. *UnTRIM San Francisco Bay–Delta Model Sea Level Rise Scenario Modeling Report*. Delta Conservation Plan. Prepared for Science Applications International Corporation and the California Department of Water Resources.

MacWilliams, M.L., E.S. Gross, J.F. DeGeorge, and R.R. Rachielle, 2007. *Three-Dimensional Hydrodynamic Modeling of the San Francisco Estuary on an Unstructured Grid*. 32nd World Congress of the International Association for Hydro-Environment Engineering and Research (Venice Italy); July 2007.

MacWilliams, M.L., F.G. Salcedo, and E.S. Gross, 2008. *San Francisco Bay–Delta UnTRIM Model Calibration Report*. POD 3-D Particle Tracking Modeling Study. Prepared for the California Department of Water Resources.

MacWilliams, M.L., A.J. Bever, E.S. Gross, G.S. Ketefian, and W.J. Kimmerer, 2015. "Three-Dimensional Modeling of Hydrodynamics and Salinity in the San Francisco Estuary: An Evaluation of Model Accuracy, X2, and the Low–Salinity Zone." *San Francisco Estuary and Watershed Science* 13(1). April 2015. Available at: <https://doi.org/10.15447/sfews.2015v13iss1art2>.

References (Continued)

MacWilliams, M.L., A.J. Bever, and E. Foresman, 2016a. "3-D Simulations of the San Francisco Estuary with Subgrid Bathymetry to Explore Long-Term Trends in Salinity Distribution and Fish Abundance." *San Francisco Estuary and Watershed Science* 14(2). Available at: <https://doi.org/10.15447/sfews.2016v14iss2art3>.

MacWilliams, M.L., E.S. Ateljevich, S.G. Monismith, and C. Enright, 2016b. "An Overview of Multi-Dimensional Models of the Sacramento–San Joaquin Delta." *San Francisco Estuary and Watershed Science* 14(4). Available at: <https://doi.org/10.15447/sfews.2016v14iss4art2>.

MacWilliams, M.L., E.S. Ateljevich, and S.G. Monismith, 2022. "How Can We Use Computer Models to Learn About the San Francisco Estuary?" *Frontiers for Young Minds* 10. March 15, 2022. Available at: <https://doi.org/10.3389/frym.2022.611920>.

Sommer, T., R. Hartman, M. Koller, M. Koohafkan, J.L. Conrad, M.L. MacWilliams, A.J. Bever, C. Burdi, A. Hennessy, and M. Beakes, 2020. "Evaluation of a Large-Scale Flow Manipulation to the Upper San Francisco Estuary: Response of Habitat Conditions for an Endangered Native Fish." *PLOS ONE* 15(10). October 1, 2020. Available at: <https://doi.org/10.1371/journal.pone.0234673>.